



Remove of lead from the polluted water by phytoremediation technique (Eclipta Alba plant)

Mohamad J. Noufal¹, Zena A. Maalla¹, Dareen J. Noufal², Ali A. Hossean³

¹Faculty of Urban Construction and Environmental Engineering, Chongqing University, 174 Shazhengjie, Shapingba District, 400045, Chongqing City, China.

² Faculty of Civil Engineering, Al-Baath University, Homs City, Syrian Arab Republic.

³ Higher Institute for Environmental Research, Tishreen University, Latakia City, Syrian Arab Republic.

Received 19 May 2016; Received in revised form 17 July 2016; Accepted 24 Aug. 2016; Available online 1 Jan. 2017

Abstract

One of the major environmental problems is the pollution of water and soils by toxic heavy metals. The aim of current study was estimation the ability of Eclipta alba plant for the removal of (Pb) from the polluted water by phytoremediation technique and study the effect of a number of parameters (Pb concentration, contact time and pH values). The plant showed higher removal efficiency (99.2%) after 7 days of treatment at 50 ppm. The best removal efficiency of Pb 98.95% at 100 ppm was occurred at pH7. The root ability to accumulate Pb was more than the shoot ability, after 7 days treatment of water and pH=7, they were 2852.5 mg/kg, 2497.5 mg/kg and 502.5 mg/kg in roots, stems and leaves respectively. These results showed that Eclipta Alba can be used in biological treatment of polluted water.

Copyright © 2017 International Energy and Environment Foundation - All rights reserved.

Keywords: Eclipta Alba; Phytoremediation; Lead; Industrial waste water.

1. Introduction

Several methods are already being used to clean up the environment from those kinds of contaminants, but most of them are costly and far away from their optimum performance. For at least 300 years, the ability of plants to remove contaminants from the environment has been recognized and taken advantage of in applications such as land farming of waste. Over time, this use of plants has evolved to the construction of treatment wetlands or even the planting of trees to counteract air, soil and water pollution [1]. Presently, phytoremediation has become an effective and inexpensive technological solution used to extract or remove inactive metals and metal pollutants from contaminated soil and water. This technology is environmental friendly and potentially cost effective. Phytoremediation takes the advantage of the unique and selective uptake capabilities of plant root systems, together with the translocation, bioaccumulation, and contaminant degradation abilities of the entire plant body [2].

Many species of plants have been successful in absorbing contaminants such as lead, cadmium, chromium, arsenic, and various radionuclides from soils and water. Among the existing pollutants, lead (Pb) is the major contaminant of the soil and water [3] which has significant environmental problems [4], including the risk of poisoning for humans and especially children [5].

Plants of the families Brassicaceae, Euphorbiaceae, Asteraceae, Lamiaceae and Scrophulariaceae have been identified as for the extraction of lead from the soil [1]. Boonyapookana et al. [6] studied the hyperaccumulation potential for Pb of sunflower, tobacco and vetiver, and concluded that all three accumulated Pb in their leaves and stems, but the first one was the most efficient accumulator. Thus, this work is aimed to study the ability of Eclipta Alba plant for lead removal from the polluted water by phytoremediation technique.

2. Materials and methods

2.1 Collection and growing of plant samples

collected plants were washed with clean tap water to remove dirt on the herbs Cuttings with leaves of about 15cm were placed in distilled water and kept for rooting in water and then transferred to beakers filled with Hoagland's hydroponics solution (containing macro and micro elements necessary for the plant growth and rooting) which prepared as described by [7].

2.2 Estimation of lead concentration in the plant tissues

The concentration of lead ion (Pb⁺²) in the plant samples (E.alba) was estimated before and after the treatment with industrial waste water.

2.3 Preparation of standard aqueous solutions

Lead standard aqueous solution was prepared depending on the atomic and molecular weight of Pb [8] as following: A standard solution of (1000 ppm) of lead Pb was prepared by dissolving 1.56g of lead acetate (Pb (C₂H₃O₂)₂) in 1000 ml of distilled water .This solution was passed through 0.45 Mm membrane filter and served as a stock solution for the preparation of the required concentrations for the experiment.

2.4 Study the efficiency of E.alba for Lead removal

2.4.1 The effect of lead concentration and contact time

Five concentrations of lead (40, 50, 65, 85,100 ppm) were selected and prepared according to the concentration of lead in the industrial waste water. Plastic containers of (40x30x15) cm was used and filled with 4 liter of each concentration. Ten well rooted propagules were put in each container and three replicates were used for each treatment. Collection of water samples from all replicate containers every day from 1 to 11 days to estimate the residual quantity of lead by Atomic absorption spectrophotometer after treatment.

2.4.2 The effect of pH

The effect of pH on the uptake of lead (Pb) by E. Alba was done by growing the plant in a range of pH values (4- 9), acidic and alkaline pHs were adjusted by adding dilute HCl and KOH respectively.

2.5 Plant analysis for estimation of pb concentration in plant tissues after phytoremediation process

For metal analysis, 0.5 g of roots stems and leaves samples were taken. 5 ml of nitric acid (65%) and 1 ml of perchloric acid (70-72%) was added [9]. The digestion was allowed to proceed in microwave digester at 80 °C for 15 minutes. It was cooled and then the digests were filled up to 50 ml with distilled water [10]. Finally, 20 ml of each sample solution was taken and used to determine Pb concentration using flame atomic absorption spectrophotometer.

2.6 Calculation of Removal efficiency (R.E)

The removal efficiency was calculated according to the following equation [11]:

$$\text{Removal efficiency (Re) \%} = \frac{C_0 - C_f}{C_0} * 100 \quad (1)$$

where: C₀ The initial metal concentration (mg/l), C_f The final concentration (mg/l)

2.7 Statistical analysis

The data were analyzed statistically by using statistical program, Anova test.

3. Result and discussion

3.1 The concentration of lead ion (Pb) in *E. Alba* tissues before phytoremediation process

The analysis to different tissues of *E. Alba* parts (roots, stems and leaves) for determination of Pb concentration in the tissues before using in treatment of polluted waste water (control) showed that the concentration of lead was 0 mg/Kg. This means that the plant is free of Pb ion in its tissue before phytoremediation process.

3.2. The removal efficiency of lead by *E. alba* from aqueous solutions

3.2.1 The effect of Pb ion concentration and time

E. Alba plant shows higher efficiency for the removal of Pb ions from aqueous solution. The highest removal efficiency (RE) was 99.2% at 50 ppm after 7 days of treatment while the lowest RE was 4.88% at 100 ppm after one day of treatment (Figure 1).




Figure 1. Effect of concentration and time on removal efficiency by *E. Alba*.

The removal efficiency (RE) was ranged from 98.28% to 4.88% (mean 67.41%) after one day of treatment. It was noted that the RE was increased with increasing of the incubation time; they reached maximum 99.2% (mean 84.65%) after seven days of incubation for all treatments. Further increase in incubation time, the RE was slowly decreased in all treatments (Table 1).

Table 1. Removal efficiency of lead at 5 different concentrations.

day	R.E% 40ppm	R.E% 50ppm	R.E% 65ppm	R.E% 85ppm	R.E% 100ppm	Mean
1	95.19	43.6	95.13	98.28	4.88	67.41
2	96.93	56.74	96.21	98.43	12.5	72.16
3	97.12	67	98.01	98.51	14.2	74.96
4	97.27	81.99	97.56	98.53	19.9	79.05
5	97.29	97.52	98.06	98.58	22.82	82.85
6	97.28	97.99	98.46	98.6	26.86	83.83
7	97.36	99.2	98.85	98.66	29.2	84.65
8	97.02	98.2	98.87	98.58	19.6	82.43
9	96.97	97.54	98.07	98.51	17.8	81.77
10	96.87	95.4	97.56	98.35	17.7	81.17
11	96.37	95	96	98.29	9.5	79.03
Mean	96.87	84.56	97.51	98.48	22.58	

The results of Anova test for the Pb remaining in the solution (Table 2) at probability level ($P \leq 0.05$) showed significant differences at the 2nd day for the concentrations (40, 50, 65, 85 ppm) while the concentration 100 ppm showed not significant differences with the other concentrations of the experiment.

The 3rd, 5th, 6th, 7th and 8th days revealed significant differences for all the concentrations (40, 50, 65, 85 and 100 ppm). At the 4th, 9th, 10th and 11th days, the concentration 85 ppm showed not significant differences with the other concentrations (40, 50, 65, and 100 ppm) which they showed significant differences at $P \leq 0.05$.

Anova test was done for each concentration of Pb with the days of the experiment and there were found that the concentrations 40, 50 and 65 ppm showed significant differences at $P \leq 0.05$ between all the 11 day of the experiment.

The concentration of 85 ppm showed not significant differences at the 4th, 9th, 10th and 11th days while showed significant differences at the 2nd, 3rd, 5th, 6th, 7th and 8th day of the experiment. On the other hand, the concentration 100 ppm showed not significant differences between the 2nd and the 3rd day while showed significant differences for all the other days of the experiment.

The metal uptake mechanism is particularly dependent on the heavy metal concentration. In the present study *E.alba* showed higher removal efficiency for Pb ion ranged from 95.19% to 98.28% at the concentrations (40, 50, 65, 85 ppm); while the lowest RE was (4.88%) at 100 ppm after the first day of treatment. In our study, the optimum Pb concentration was 50 ppm because it gives the highest removal efficiency (99.2%) after 7 days of treatment (Figure 1).

It is clear from Figure 1 that the removal efficiency of Pb ion increases at lower and medium concentrations while decreased at higher concentrations. These observations can be explained by the fact that at low concentrations of metal ion (Pb), the ratio of sorptive surface area to the total metal ions available is high and thus, there is a greater chance for Pb removal capacity. When Pb ion concentrations are increased, binding sites become more quickly saturated, therefore, more Pb ions were left unabsorbed in solution at higher concentration levels.

Contact time is one of the impact parameter on the successful of phytoremediation process. It is evident from Figure 2 that the removal of metal ion increases as the contact time increases initially, the rate of uptake is fast. However, equilibrium was attained in 7 days. Further increase in contact time to 11 days, the percentage adsorption decreases very slowly. Therefore, the contact time was optimal at 7days. Similar data was recorded by [12] when he used Duckweed (*Lemna minor*) in the removal of some heavy metals. Gallardo et al.[13] found that after one week of exposure of *Hydrilaverticillata* to concentrated lead solution, maximum uptake (98%) of Pb was observed. Al- Bayati [14] found that the RE of Pb by *Eichhorniacrassipes* reached to (100%, 100%, 99.6%, 99.6%, 99.4%, 99.4%, 98.9%, and 96.33%) for the concentrations (1, 3, 6, 10, 12, 15, 20 and 30 ppm) respectively after 10 days of treatment

3.2.2 The effect of pH on the removal of lead

The pH value of the solution is an important controlling parameter in the removal of heavy metals. It seems clearly that increasing the pH values have significant effects up on the ability of *E .alba* on removal of Pb ions.

Table 2. The residual lead remaining after treatment by *E. Alba* at different concentrations of Pb.

Day	Residual Pb at 40ppm	Residual Pb at 50ppm	Residual Pb at 65ppm	Residual Pb at 85ppm	Residual Pb at 100ppm
1	1.92	28.2	3.16	1.45	95.12
2	1.22	21.63	2.46	1.33	87.5
3	1.15	16.5	1.28	1.26	85.8
4	1.08	9.002	1.58	1.24	80.1
5	1.08	1.24	1.25	1.20	77.18
6	1.08	1.004	0.99	1.17	73.14
7	1.05	0.4	0.74	1.13	70.8
8	1.19	0.9	0.79	1.20	80.4
9	1.21	1.23	1.25	1.26	82.2
10	1.25	2.3	1.58	1.40	82.3
11	1.45	2.5	2.6	1.45	90.5




Figure 2. The removal efficiency of Pb at 6 different pH values.

The results were presented in Table 3, showed that the removal efficiency (RE) of Pb ions from aqueous solution at (100 ppm) was affected by the pH. The higher RE reached 100% was observed at pH 9 and after 7 days of treatment, while the minimum RE was 0% at pH 4 after one day of treatment. It was noticed that as the pH increases, the RE also increases and reaches to the optimum values at pH above 6.

Table 3. The removal efficiency of Pb by *E. Alba* after using 6 different pH values.

Day	R.E% pH=4	R.E% pH=5	R.E% pH=6	R.E% pH=7	R.E% pH=8	R.E% pH=9
1	0	12.4	4.88	49.6	72.6	91.5
2	3.7	12.8	12.5	63.2	76.6	93.1
3	4.5	18.5	14.2	95.1	97.2	99.15
4	5.1	20.7	19.9	95.2	97.48	99.85
5	6.5	21.2	22.82	98.28	97.93	99.61
6	7.8	51.6	26.86	98.91	98.21	99.89
7	9.5	52	29.2	98.95	98.73	100
8	6.2	32.3	19.6	98.88	99.11	100
9	1.7	26.4	17.8	98.32	98.2	100
10	0	22.7	17.7	97.61	98.21	100
11	0	18.84	9.5	97.19	98.25	100

The results also showed that with pH values 8 and 9 (above 7), Pb ions are precipitated as lead hydroxide [15] and this case could give wrong initial concentration. Therefore, the optimum pH for removal of Pb ions fixed as pH 7 because it gives the highest RE (98.95%) after 7 days of treatment Figure 2.

In addition, it was interesting to note that there was fluctuation of pH values for Pb solution after application of *E. Alba*. Both acidic and alkaline pH of the solution shifted slightly towards neutral pH with increase in the number of days. This change in pH may be due to release of some root exudates in response to the stress so as to adapt itself to the existing environment. Thus, the result of present study shows that the plant *E. Alba* has the pH neutralizing capacity.

A nova test for the remaining of Pb ions in the solution (Table 4) revealed significant effects of increasing pH values at levels ($P \leq 0.05$), also this metal was significantly affected by these pH levels at the same probability.

It was noticed that the ability of removing lead ions depends on the pH values of the solution and this depends on the ion state and nature of material. At low pH value, binding sites are generally protonated

or positively charged (by the hydronium ions). Thus, repulsion occurs between the metal cation and the adsorbent. As the pH value increase, the binding sites start deprotonating and makes different functional groups available for metal binding. In general, cation binding increases as pH increases [16, 17].

The result obtained from current study was in accordance with the results of other authors. Goswami *et al.* [18] found that *Eichhorniacrassipes* has the maximum removal efficiency of Pb at neutral pH.

Baharudin and Mohd [19] found that the constructed wetland containing 15-plants recorded the highest removal with 99.28% for lead at neutral condition (pH 7) compared to the base and acidic conditions.

However other studies obtained different results either less or more than of our study (pH 7). The amount of lead absorbed by plants is affected by the pH and to reduce lead uptake by plants, the pH of the soil or water is adjusted to a level of 6.5 to 7 [20].

While, Uysal and Taner [21] examined the ability of the *L. minor* to remove soluble lead under different pH values (4.5-8.0) and temperature (15-35°C) in presence of different Pb concentrations 0.1-10.0 mg/L for 7 days. They notice Pb accumulation at pH 4.5 and then it decreased to pH 6, but it did not change at pH range 6-8.

Table 4. The residual Pb ion remaining in the solution after treatment by *E. Alba* at different pH values.

Day	pH=4	pH=5	pH=6	pH=7	pH=8	pH=9
1	100	87.6	95.12	50.4	27.4	8.5
2	96.3	87.2	87.5	36.8	23.4	6.9
3	95.5	81.5	85.8	4.9	2.8	0.85
4	94.9	79.3	80.1	4.8	2.52	0.15
5	93.5	78.8	77.18	1.72	2.07	0.39
6	92.2	48.4	73.14	1.09	1.79	0.11
7	90.5	48.03	70.8	1.05	1.27	0
8	93.8	67.7	80.4	1.12	0.89	0
9	98.3	73.6	82.2	1.68	1.80	0
10	100	77.3	82.3	2.39	1.79	0
11	100	81.16	90.5	2.81	1.75	0

3.3 The accumulated lead (Pb) ion in *E. Alba* tissues after phytoremediation process

According to plant analysis with dry weight, the concentration of Pb in *E. Alba* tissues after 7 days at pH 7 was estimated. The average values 2852.5, 2497.5 and 502.5 mg/kg were observed in roots, stems and leaves respectively (Figures 3 and 4). The results showed that *E. Alba* had the ability to accumulate large quantities of lead, especially in roots comparing to stems and leaves. Ashokkumar *et al.* [22] reported that *E. Alba* has the ability to uptake and accumulate heavy metals including Pb from polluted soils.

Metal concentrations in plants vary with plant species [23, 24, 25 and 26]. The high metal uptake may be attributed to highly efficient intracellular compartmentalization. Effective parameters of metal uptake by plants were reviewed to be genetic architecture of metal accumulation as well as binding ligands such as organic acids, amino acids, peptides and proteins including citrate, malate, histidine, nicotineamine, glutathione, phytochelatins, metallothioneins [27].




Figure 3. Lead concentrations (mg/kg) in *E. Alba* parts after phytoremediation process.



Figure 4. E. Alba plant in the polluted water after 7 days of incubation at pH 7.13.

Acknowledgements

This work was supported by International Energy and Environment Foundation.

References

- [1] United States Environmental Protection Agency (USEPA). 2000. Introduction to Phytoremediation. EPA 600/R-99/107, U.S. Environmental Protection Agency, Office of Research and Development, Cincinnati, OH.
- [2] Hinchman, R. R.; Negri, M. C. and Gatliff, E. G. 1995. "Phytoremediation: using green plants to clean up contaminated soil, groundwater, and wastewater," Argonne National Laboratory Hinchman, Applied Natural Sciences, Inc.
- [3] Gratão, P.L.; Prasad, M.N.V.; Cardoso, P.F.; Lea, P.J. and Azevedo, R.A. 2005. Phytoremediation: Green technology for the cleanup of toxic metals in the environment. *Braz. J. Plant Physiol.* 17:53-64.
- [4] Shen, Z.G.; Li, X.D.; Wang, C.C.; Chen, H.M. and Chua, H. 2002. Lead phytoextraction from contaminated soil with high-biomass plant species. *J. Environ. Qual.* 31:1893-1900.
- [5] Lasat, M. M. 2000. "Phytoextraction of metals from contaminated soil: a review of plant/soil/metal interaction and assessment of pertinent agronomic issues," *J. of Hazardous Sub. Res.*, 2(5): 1-25.
- [6] Boonyapookana, B.; Parkplan, P.; Techapinyawat, S.; De Laune, R.D. and Jugsujinda, A. 2005. Phytoaccumulation of lead by sunflower (*Helianthus annuus*), tobacco (*Nicotianatabacum*), and vetiver (*Vetiveriazizanioides*). *J. Environ. Sci. Health Part A-Toxic/Hazard. Subst. Environ. Engin.* 40:117-137.
- [7] Jasoni, R.L.; Cothren, J.T.; Morgan, P.W. and Sohan, D.E. 2002. Circadian ethylene production in cotton. *Plant Growth Regulation*, 36(2): 31-37.
- [8] Harley, J.P. and Prescott, L.M. 1996. *Laboratory exercises in microbiology*. 3rd ed. McGraw-Hill, Boston. USA.
- [9] Wang, X. and Zhou, Q.X. 2003. Distribution forms of cadmium, lead, copper and zinc in soil and its influences by modifier. *J. Agro-Environ. Sci.*, 22: 541-545.
- [10] Price, W.J. 1979. *Spectrochemical Analysis by Atomic Absorption*. Heydon and Sons Ltd. London. 254-255.
- [11] Fahad, H. G. 1994. A study of efficiency of different microorganisms in thorium sorption from aqueous solutions. M.Sc. thesis. College of Science, Baghdad University, Iraq.
- [12] Dhabab, J. M. 2011. Removal of some heavy metal ions from their aqueous solutions by duckweed. *J. Toxicol. Environ. Health Sci.*, 3(6): 164-170.
- [13] Gallardo, T.; Maria, B., Robert, F. and Martin F. 1999. Lead accumulation by three aquatic plants. *Symposia papers presented before the division of Environmental Chemistry, American Chemical Society*, 39(2): 46-47.
- [14] Al-Bayati, S. M.H. 2008. Removal of Copper and Lead metals from water ecosystem by Water hyacinth *Eichhorniacrassipes*(Mart.) Solms. M.Sc. Thesis, College of Science for women, University of Baghdad, Iraq.
- [15] Lindsay, W. L. 1979. *Chemical equilibria in soils*. John Wiley and Sons. New York.
- [16] Forsner, U. and Wittman, G.T. 1981. *Metal pollution in the aquatic environment*. Springer-Verlag, Berlin, Germany, 2nd edition.
- [17] Esposito, A.; Pagnanelli, F. and Veglio, F.I. 2002. Plant proving their worth in toxic metal. *Chem. Eng. Sci.*, 57: 307-313.

- [18] Goswami, R.; Thakur, R. and Sarma, K.P.2010 .Uptake of Lead from Aqueous Solution using Eichhorniacrassipes: Effect on Chlorophyll Content and Photosynthetic Rate.Int.J. Chem.Tech. Res., 2(3):1702-1705.
- [19] Baharudin, B. & Mohd, S. 2008. Lead and cadmium removal in synthetic wastewater using constructed wetland. Faculty of Chemical & Natural Resources Eng. University, Malaysia Pahang.
- [20] Traunfeld, J. H. and Clement, D. L.2001. "Lead in Garden Soils. Home and Garden," Maryland Cooperative Extension, University of Maryland.
- [21] Uysal, Y. and Taner, F. 2009. Effect of pH, Temperature and Lead Concentration on the Bioremoval of Lead from water using *L. minor*. Int. J. Phytoremed., 11: 591-608.
- [22] Ashokkumar, B.; Jothiramalingam, S.; Thiyagarajan, S.K., Hidhayathullakhan, T. and Nalini, R. 2014. Phytoremediation of tannery polluted soil using *Eclipta Alba* (karisalankanni). Int.J.Curr.Res. Chem.Pharma.Sci. 1(3):01-05.
- [23] Alloway, B.J.; Jackson, A.P. and Morgan, H. 1990. The accumulation of cadmium by vegetables grown on soils contaminated from a variety of sources, Science Total Environment, 91:36- 223.
- [24] Secu, C.V.; Iancu, O.G. and Buzgar, N. 2008. Lead, zinc and copper in the bioaccumulative horizon of soils from Iași and the surrounding areas, Carpathian J. of Earth and Environ. Sci., 3(2):131-144.
- [25] Petrescu, L. and Bilal, E. 2007. Environmental impact assessment of a uranium mine, east Carpathians, Romania: metal distribution and partitioning of U and Th, Carpathian J. of Earth and Environ. Sci., 2(1): 39-50.
- [26] Lăcătușu, R.; Cîtu, G; Aston, J.; Lungu, M. and Lăcătușu, A.-R. 2009. Heavy metals soil pollution state in relation to potential future mining activities in the RoșiaMontană area, Carpathian J. of Earth and Environ. Sci., 4(2): 39-50
- [27] Verbruggen, N.; Hermans, C. and Schat, H. 2009. Molecular mechanisms of metal hyperaccumulation in plants. New Phytol., 181(4):759-776.



Mohamad J. Noufal is PhD student in College of Urban Construction and Environmental Engineering, Chongqing University. He has got his master of science in Environmental Engineering from Chongqing University. He has got his bachelor of science in Environmental and civil Engineering from Al-Baath University in Syria. He also has four years of work experience in the Industrial wastewater treatment in Syria. At present, his research interests mainly focus on wastewater management system.
E-mail address: mohmadnofal@hotmail.com



Zena A. Maalla is master student in College of Urban Construction and Environmental Engineering, Chongqing University. She has got his bachelor of science in Environmental and civil Engineering from Al-Baath University in Syria. She also has three years of work experience in Water treatment plants in Syria.
E-mail address: zenamaalla1990@hotmail.com



Dareen J. Noufal has got her Master degree of science in civil Engineering from Al-Baath University in Syria.



Ali A. Hossean is master student in Higher institute for environmental Research, Tishreen University. He has got his bachelor of science in Environmental and civil Engineering from Al-Baath University in Syria
E-mail address: ali.hossean.1985@gmail.com